2011 WSU Variety Testing Soft White Spring Wheat Trial Summary Precipitation Zone 12-16"

Variety Name (Club Italicized)	Almira	Endicott	E Lamont	Average		Almıra	Endicott	다 디	Average	Almira	Endicott	Lamont	
_ouise-G2	71	61	89	74		9.7	62.3	61.2	61.1	10.5	9.9	10.1	
Diva	80	56	81	72		0.6	61.5	61.5	61.2	9.8	10.3	9.5	
Louise	68	56	88	71		0.2	62.3	61.6	61.4	10.3	10.1	9.6	
WA 8131	78	54	81	71		0.4	62.3	61.5	61.4	10.9	10.6	9.9	
JD	75	57	78	70		1.2	63.6	62.7	62.5	10.9	10.2	10.4	
DO687	76	51	77	68	6	2.0	63.7	62.3	62.7	10.2	10.0	9.4	
Babe	68	54	83	68	6	1.5	63.0	61.3	61.9	10.1	9.9	9.8	
DO686	73	49	78	67	6	0.1	63.5	62.4	62.0	10.0	10.5	9.8	
Whit	69	55	78	67	6	0.9	62.3	60.8	61.3	10.4	10.0	10.1	
DO671	72	48	81	67	6	0.0	62.6	61.3	61.3	9.7	10.0	9.3	
WA 8150	70	55	74	66	6	1.3	63.5	62.0	62.2	9.9	10.1	9.8	
WA 8128	71	50	76	66	6	1.5	62.7	62.0	62.1	10.8	10.2	10.7	
WA 8127	71	51	75	66	6	0.9	62.9	61.6	61.8	10.0	10.2	9.7	
Alturas	68	49	80	66	5	8.9	62.2	60.7	60.6	10.6	9.9	9.4	
Eden	69	55	73	66	6	0.7	63.2	62.5	62.2	10.1	9.9	9.9	
WA 8149	69	56	71	65	6	1.2	61.6	60.6	61.1	9.2	10.2	9.1	
IDO644	63	54	80	65	6	0.4	61.3	60.4	60.7	10.7	9.6	8.9	
WA 8124	55	56	82	64	6	0.7	63.7	61.8	62.1	10.1	10.5	9.6	
Wakanz	61	54	76	64	6	0.1	61.0	59.7	60.3	11.2	10.4	10.5	
Zak	62	49	67	59	6	1.0	62.6	60.1	61.3	10.6	10.6	10.2	
Nick	57	51	68	59	6	0.9	61.9	60.7	61.2	9.9	10.0	9.3	
UI-Cataldo	60	46	71	59	5	9.0	61.8	60.6	60.5	10.3	10.4	10.0	
Alpowa	56	47	71	58	6	1.7	63.2	61.6	62.2	9.5	10.1	8.9	
WB-1035CL2	54	43	64	54	6	0.4	61.6	60.9	60.9	11.5	11.1	11.1	
CV (%)	13	4	6	9	1	1.8	0.6	0.6	1.2	4.4	2.1	5.9	
LSD (0.10)	9	4	5	4	1	1.2	8.0	0.4	0.4	0.5	0.4	0.6	
Average	67	52	77	65	6	0.6	62.5	61.3	61.5	10.3	10.2	9.8	
Highest	80	61	89	74	6	2.0	63.7	62.7	62.7	11.5	11.1	11.1	
Lowest	54	43	64	54	5	8.9	61.0	59.7	60.3	9.2	9.6	8.9	

2011 WSU Soft White Spring Wheat Trial Summary

Precipitation Zone 12-16" – Preliminary Data

- 1. Soft white spring wheat grain yield across three locations and 24 entries in the 12-16" precipitation zone averaged 65 bushels/acre and is five bushels/acre higher than the 2010 average of 60 bushels/acre. The CV for the average data is 9, lower than the 2010 CV. In general the trials had good establishment after late seeding.
- 2. Yields among entries averaged across locations ranged from 54 to 74 bushels/acre. Louise-G2 was the highest yielding entry averaged across locations. The Louise-G2 entry is a 2 oz/100lb of seed treatment of Gaucho. This treatment targets wire worms that may be causing yield losses and appears to be effective at the Almira and Endicott sites. Average yield values within the 10% LSD range (4 bushels/acre) of the highest yield are shown in bold and this included 5 of the 24 entries. Stripe rust was a factor at all locations and influenced yield rankings based on susceptibility. There were no fungicide applications made on these trials and stripe rust yield impacts in percent are: 20% impact at Almira, 10% at Endicott, and 20% at Lamont.

 3. Test weight averaged 61.5 lb/bu across locations and entries and was higher than last year's 58.1 lb/bu
- 3. Test weight averaged 61.5 lb/bu across locations and entries and was higher than last year's 58.1 lb/bu average. Grain protein averaged 10.1% and was lower than last year's 13.0% protein value.