2011 WSU Variety Testing SW Spring Wheat Trial, Bickleton

Variety Name *Club Italized	5 YEAR AVERAGE (BU/A)	3 YEAR AVERAGE (BU/A)	2 YEAR AVERAGE (BU/A)	2011			
				YIELD (BU/A)	TEST WT (LBS/BU)	PROTEIN (%)	PLANT HT
JD		28	29	32	62.0	11.7	21
Wakanz		27	28	30	59.1	11.3	23
Diva		24	24	30	61.6	10.5	25
Louise-G2				28	60.9	10.7	25
UI-Cataldo		21	21	27	59.2	11.1	23
Alturas		24	23	26	60.0	11.0	23
Babe		28	29	26	61.8	10.9	23
IDO687				25	60.7	10.4	21
Alpowa		23	24	24	58.5	10.8	22
Louise		24	23	24	60.6	10.7	25
Whit		25	25	24	60.9	10.5	21
Zak		25	24	23	61.3	12.1	21
IDO644				23	58.9	10.9	22
WA 8131				22	59.4	11.1	21
Eden		25	25	21	61.7	11.4	18
WA 8128				21	61.6	11.6	20
WA 8149				20	60.5	11.4	19
WA 8150				20	60.6	10.5	22
WA 8124			21	19	60.3	10.8	23
IDO686				19	61.2	10.9	25
WA 8127				18	60.8	11.6	20
IDO671			19	17	59.7	10.5	19
WB-1035CL2				17	58.5	12.0	21
Nick		25	24	16	59.8	11.3	19
C.V.		29	28	29	1.8	4.9	12
LSD		3	4	N.S. ¹	1.2	0.6	3
Average		25	24	23	60.4	11.1	22
Highest		28	29	32	62.0	12.1	25
Lowest		21	19	16	58.5	10.4	18

^{1 -}No Significant Difference

Bickleton Soft White Spring Wheat – Preliminary Data

- 1. Grain yield in the Bickleton soft white spring wheat trial averaged 23 bushels/acre 2 bushels/acre less than the 3-year average. The Bickleton nursery was located about four miles east of Bickleton, WA (Steve Matsen, cooperator).
- 2. This nursery was seeded on 12 May, 2011 following spring wheat. Seed was placed at a 60#/acre seeding rate using a no-till plot drill set on 10-inch spacing. Base fertilizer was 30 #N/acre applied at seeding. Seeding was late and emergence was slow due to cool temperatures.
- 3. Yields ranged from 16 bu/acre to 32 bu/acre. Due to low yields and high variability at this site, yield was not significantly different among entries. JD and Babe were highest yielding across 3 years at this location. Stripe rust was a minor factor in this trial and no fungicide was applied.
- 4. Test weights were good and averaged 60.4 lb/bu and ranged from 58.5 to 62.0 lb/bu. Grain protein was good and averaged 11.1% with a range of 10.4 to 12.1/%. The average plant height was 22 inches with no lodging.