

W.J. Johnston<sup>1</sup>, R.C. Johnson<sup>2</sup>, K.L. Dodson<sup>1</sup>, G.K. Stahnke<sup>1</sup>, and C.T. Golob<sup>1</sup>

<sup>1</sup>Department of Crop and Soil Sciences, Washington State University, Pullman, WA

<sup>2</sup>Western Regional Plant Introduction Station, Pullman, WA

## INTRODUCTION

A ban on open-field burning of Kentucky bluegrass (*Poa pratensis* L.) post-harvest residue has been implemented in Washington and restrictions are in place in Idaho and Oregon, USA. Without post-harvest burning seed yield decreases over time (Lamb and Murray, 1999), which has forced growers to use shorter rotations to maintain yield. In a multi-year study we have identified germplasm that has improved seed production without burning (Johnson et al., 2003), reselected within that germplasm (Johnson et al., 2010), increased seed, and established turfgrass and seed production trials (Dodson, 2008). These trials will be evaluated over several years.

## OBJECTIVES

Develop a high yielding turf-type Kentucky bluegrass that that can be grown for several years without open-field burning of post-harvest residue.

## MATERIALS AND METHODS

This long-term study initially evaluated 228 bluegrasses from the USDA-ARS Kentucky bluegrass collection at Pullman, WA (Fig. 1; Johnston et al., 1997). The selected eight PI accessions (plus two checks, 'Kenblue' and 'Midnight') represent germplasm that has good seed yield without field burning of post-harvest residue while maintaining turfgrass quality (Fig. 2 and 3). Agronomic yield parameters were evaluated over a 2-yr period and individual plants were selected within each accession, or check, with the highest seed weight, highest seed panicle<sup>-1</sup>, highest panicles area<sup>-1</sup>, and highest seed yield (Fig 4). These 40 selections plus remnant seed obtained from the USDA-ARS collection (base population) were planted in turf plots in 2006 (Fig. 5) and non-irrigated and irrigated seed production plots in 2007 at Pullman, WA (Fig. 6). There were 150 plots in each trial (50 entries x 3 replications). The turfgrass trial was evaluated monthly during the growing season (2007 to 2010) according to National Turfgrass Evaluation Program (NTEP) protocol. From 2008 to 2010, seed production plots were harvested, threshed, cleaned, and seed yield was determined. Only results for 2010 are presented (3<sup>rd</sup> harvest).



Fig. 1 USDA-ARS Kentucky bluegrass collection evaluation



Fig. 2 Seed plot treatments: burning; baling; full residue



Fig. 3 Turfgrass evaluation: turf quality; texture; color; etc.



Fig. 4 Space-plant nursery for individual plant characterization



Fig. 5 Turfgrass evaluation at Pullman, WA



Fig. 6 Seed production evaluation at Pullman, WA




Fig. 7 Non-irrigated Kentucky bluegrass seed yield vs. turf quality (1 to 9; 9 = excellent quality) at Pullman, WA 2010.




Fig. 8. Irrigated Kentucky bluegrass seed yield vs. turf quality (1 to 9; 9 = excellent quality) at Pullman, WA 2010.

## RESULTS

### Evaluation of the USDA-ARS Kentucky Bluegrass Collection

- Agronomic data could be used to differentiate among accessions.
- A Kentucky bluegrass core was developed (Johnston et al., 1997).

### Residue Management and Turf Evaluation

- Accessions were identified that maintained good seed yield when post-harvest residue was baled and possessed good turf quality (Johnson et al., 2003).

### Selection for Diversity in Seed Yield Components

- Variation between and within accessions was identified (Johnson et al., 2010).

### Turfgrass and Seed Production Trials

- A turfgrass evaluation trial was established at Pullman, WA (fall 2006). Plots were evaluated for turfgrass parameters, 2007 to 2010.
- Seed production trials (irrigated and non-irrigated) were established at Pullman, WA in 2007 and harvested 2008 to 2010.

## OUTCOMES & CONCLUSIONS

- The USDA-ARS Kentucky bluegrass collection was evaluated for diversity and a core collection was developed.
- Accessions with good seed yield without post-harvest residue burning and good turfgrass quality were identified.
- Variation within accessions for seed production parameters was found, so the potential exists for plant selection and enhancement.
- Seed production and turfgrass trials were established and evaluations are on-going (2010 was 3<sup>rd</sup> of four, or more, harvests).
- Selection for seed yield components had a variable response.
- PI 368241 and Kenblue selection for seed panicle<sup>-1</sup> (non-irrigated) show promise of being able to provide good turfgrass quality and seed yield under non-burn residue management (Fig. 7 and 8).

## LITERATURE CITED

- Dodson, K.L. 2008. Development of Kentucky bluegrass (*Poa pratensis* L.) for non-burn seed production. M.S. thesis. Washington State Univ., Pullman.
- Johnson, R.C., W.J. Johnston, and C.T. Golob. 2003. Residue management, seed production, crop development, and turf quality in diverse Kentucky bluegrass germplasm. *Crop Sci.* 43:1091-1099.
- Johnson, R.C., W.J. Johnston, F.B. Bertoli, and C.T. Golob. 2010. Seed yield, development, and variation in diverse *Poa pratensis* accessions. *Crop Sci.* 50:337-344.
- Johnston, W.J., M.C. Nelson, R.C. Johnson, and C.T. Golob. 1997. Phenotypic evaluation of Kentucky bluegrass (*Poa pratensis* L.): USDA/ARS Plant Introduction germplasm collection. *Int. Turf. Soc. Res. J.* 8:305-311.
- Lamb, P.F., and G.A. Murray. 1999. Kentucky bluegrass seed and vegetative responses to residue management and fall nitrogen. *Crop Sci.* 39:1416-1423.

## ACKNOWLEDGMENTS

Research was partially funded by a GSCSSA USDA-ARS special grant, Washington State Department of Ecology, and Washington Turfgrass Seed Commission.

ASA-CSSA-SSSA San Antonio, TX 2011