

Tree Pests and Pathogens of Concern



Field Guide

treehealth.wsu.edu

FIELD GUIDE

Purpose

The purpose of this guide is to provide information to help monitor for the accidental introduction of novel plant pests and diseases. Early detection is key to reducing the impacts of introduced species on northwest forests and communities.

Partners

This booklet was made in partnership with the U.S. Department of Agriculture (USDA) and Washington State University.

Funding

This booklet was designed by Washington State University and produced with support from the USDA Animal and Plant Health Inspection Service (APHIS), USDA Forest Service, and USDA Forest Service International Programs.

USDA is an equal opportunity provider, employer, and lender.

WSU Urban Forest Health Lab

The mission of the WSU Urban Forest Health Lab is to keep trees and communities healthy through discovery, applied learning, and engagement.

More information about the lab is available at <https://treehealth.wsu.edu/>

Table of Contents

Pests

| | |
|----------------------------------|---|
| Mediterranean Oak Borer | 1 |
| Emerald Ash Borer | 2 |
| Asian Longhorn Beetle | 3 |
| Citrus Longhorn Beetle | 4 |
| Red-Necked Longhorn Beetle | 5 |
| Spotted Lanternfly | 6 |
| Brown Marmorated Stink Bug | 7 |
| Spongy Moth | 8 |
| Elongate Hemlock Scale | 9 |

Diseases

| | |
|-------------------------------|----|
| <i>Phytophthora</i> spp. | 10 |
| Laurel Wilt | 12 |
| Ash Dieback | 13 |
| Oak Wilt | 14 |
| Sooty Bark Disease | 15 |
| Dutch Elm Disease | 16 |
| Beech Leaf Disease | 17 |

Front Cover Image Credits: Top: Joseph Hulbert, WSU; Middle: Richard Gardner, Bugwood.org; Bottom: David Cappaert, Bugwood.org

This booklet was created by F Dewitz, Z Gallardo, J Hulbert, and M Elliott at Washington State University.

WSU Ornamental Plant Pathology Program

The WSU Ornamental Plant Pathology program specializes on pests and diseases of ornamental plants and Christmas trees.

More information about the program is available at <https://ppo.puyallup.wsu.edu/>



WASHINGTON STATE
UNIVERSITY



Tree Pests and Pathogens of Concern



Adult. Image Credit: Pest and Diseases Image Library, Bugwood.org

Hosts

Oaks (*Quercus* spp.)

Identification

Cylindrical beetles that are reddish brown in color and 3.0 - 3.2 mm long and three times as long as they are wide. Adult females are larger and can fly, while males are smaller (2.3 mm), incapable of flying, and has a rhino horn-like structure. These beetles can be seen nearly all year in galleries or outside the tree.

Signs and Symptoms

- Wilting or stunted leaves
- Defoliation or flagging (browning or absent leaves on individual branches) on top third of tree
- Dead or broken branches, sometimes dying close to the trunk first
- Thinning crown
- Canopy dieback
- Black-stained galleries primarily located in the sapwood
- Frass accumulated on branches, bark, or around the base of the tree
- Perfectly round entrance and exit holes (1.6 mm)

Image Credits: Top Left & Right: Curtis Ewing, CALFIRE; Bottom Left: Bob Rabaglia, U.S. Forest Service; Bottom Right: Dr. Akif Eskalen, UC Davis



Adult Mediterranean Oak Borer



Galleries



Galleries



Canopy dieback



Emerald Ash Borer (*Agrilus planipennis*)



Adult. Image Credit: David Cappaert, Bugwood.org

ACTUAL SIZE

Hosts

Ash trees (*Fraxinus* spp.)

Signs and Symptoms

- D-Shaped Exit Holes in Bark (2.5 - 5 mm)
- Serpentine galleries under bark from larval feeding
- Unhealthy Ash Trees:
 - Branch Dieback
 - Canopy Dieback
 - Woodpecker Damage
 - Epicormic Branching

Identification

Adults: Beetles can be 8.5 - 19 mm long and are typically bright, metallic green. With its wings open, you can see their coppery, red abdomen. Emerge from early June to late August.

Larvae: Found inside the galleries, larvae can be up to an inch long. The bodies are creamy white with segmented, bell-shaped bodies. The larvae are typically inside the tree in June-October.

Image Credits: Top Left & Right: Eric R. Day, Virginia Polytechnic Institute and State University; Bottom Left: Art Wagner, USDA-APHIS; Bottom Right: Joseph OBrien, USDA Forest Service



Adult in galleries



Branch/ Canopy Dieback



Galleries



D-Shaped Exit Holes



Asian Longhorn Beetle (*Anoplophora glabripennis*)



ACTUAL
SIZE

Adult. Image Credit: Gillian Allard, FAO of United Nations

Hosts

Preferred US Hosts: Maples, birches, elms, horsechestnut, and willow (hardwoods)

Identification

Adults: 2.5 - 3.8 cm long beetles that are shiny black with irregular white spots and distinctive bluish-white legs. Adults have long black and white banded antennae. Seen late spring to fall.

Larvae: Creamy, white and grub-like, reaching up to 5 cm long. Larvae are typically inside the tree.

Signs and Symptoms

- Round dime sized exit holes
- Sap may be leaking from exit hole wounds
- Yellowing Leaves
- Branch dieback and flagging
- Premature leaf drop
- Oval shaped egg pits dug into trunks, branches, or exposed roots
- Frass around the base of the trunk

Image Credits: Top Left & Bottom Right: Joe Boggs, Ohio State University; Top Right: Pennsylvania Department of Conservation and Natural Resources - Forestry; Bottom Left: Steven Katovich



Emergence Hole



Branch Dieback



Larvae



Galleries



Citrus Longhorn Beetle (*Anoplophora chinensis*)



ACTUAL
SIZE

Adult. Image Credit: Ken-ichi Ueda, iNaturalist

Hosts (This list is not exhaustive)

Hardwoods including citrus spp., apple, alder, birch, pecan, chestnut, Japanese cedar, beech, fig, ash, mallow, holly, walnut, plane tree, poplars, cherry, peach, apricot, plum, pear, oak, sumac, rose, and elm trees

Identification

Adults: Shiny black beetles with 10 - 12 round white spots. Females are 3.5 cm long and males are 2.5 cm long. Adults have long black and white banded antennae. Seen April to August but some can be present in fall.

Larvae: Creamy white to yellowish brown with an amber head. Legless grubs up to 5.2 cm long.

Signs and Symptoms

- Large 6 - 9 mm diameter, round/oval exit holes
- Holes are deep into the tree to reach the xylem
- Weak, breaking branches on healthy-looking trees
- Thinning canopy and branch dieback
- Yellowing and drooping leaves
- Frass on the ground, on branches, or on bark
- Oozing sap
- Cracked or missing bark
- Round or oval egg pits dig into the bark, 1.3 cm in diameter
- Larval tunnels in the wood

Image Credits: Top & Bottom Left: Art Wagner, USDA APHIS; Top Right: National Plant Protection Organization, the Netherlands, Bugwood.org; Bottom Right: UK Crown Copyright Courtesy of Fera/WTML



Adult



Larvae



Emergence Holes



Adult with Exit Hole



Red-Necked Longhorn Beetle (*Aromia bungii*)



Image Credit: Gillian Allard, FAO of United Nations

Hosts

Prunus spp. (cherry, plum, apricot, peach, ornamentals)

Signs and Symptoms

- Oval shaped exit holes, 6 - 10 mm wide and 10 - 16 mm long
- Yellow to red leaves
- Wilting leaves and early leaf fall
- Galleries inside the tree
- Frass at the base of the tree trunk
- Eggs laid in bark crevices

Identification

Adults: Elongate, glossy black beetles with a red thorax between its head and abdomen. Beetles are 20 - 40 mm long. Female antennae are as long as its body and male antennae are 1 ½ times as long. Adults emerge between June and August

Larvae: Mature larvae are yellowish white. The larvae are typically inside the tree. Length ranges from 40 - 55 mm

Image Credits: Top Left: Fera Science Ltd.; Top Right, Bottom Left, & Bottom Right: Raffaele Griffo, Plant Protection Service Regione Campania, Napoli, Italy




Adult



Plant infested by *Aromia bungii* larvae



Emergence Holes



Eggs



Spotted Lanternfly (*Lycorma delicatula*)



Adult. Image Credit: Pennsylvania Department of Agriculture

Hosts

Preferred host: Tree of heaven (*Ailanthus altissima*)
The insect feeds on a wide variety of other hardwood trees and shrubs

Signs and Symptoms

- Branch Dieback
- Sap oozing from open wounds
- Wilting foliage
- Clear, sticky substance at bottom of tree called honeydew
- Black fungus growing on top of honeydew called sooty mold
- Increased ant, bee, and wasp activity from exposed honeydew and sap

Identification

- Adults: 20.5 - 26.5 mm long and roughly 12.5 mm wide with a yellow with black striped abdomen and large, visually striking wings. The forewings are brown spotted with black tips and hind wings are scarlet towards the center, white striped with black tips. Adults can be seen June - December.
- Nymphs: Early in their development appear black with white spots, seen spring - summer. Turn red with white spots before becoming adults, seen late summer - early fall
- Eggs: Laid in mass, the eggs are yellowish-brown and covered with a gray, waxy coating. Coating darkens as egg mass weathers. Found on smooth surfaces September to June.

Image Credits: Top Left: Colin Purrington; Top Right: Lawrence Barringer, Pennsylvania Department of Agriculture; Bottom Left: Richard Gardner, Bugwood.org; Bottom Right: Emelie Swackhamer, Penn State University



Early stage nymph



Mature nymph



Adults with wings closed



Eggs



Brown Marmorated Stink Bug (*Halyomorpha halys*)



Adult. Image Credit: Susan Ellis, Bugwood.org



Hosts (this list is not exhaustive)

Preferred hosts: apple, pear, peach, grape, blueberry, soybean, tomato, corn, ornamental trees and shrubs

Signs and Symptoms

- Damaged fruits
- Discolored, irregular depressions on fruit surface
- Necrotic spots
- Misshapen fruit
- Early fruit drop
- Stippled yellow, white, or brown areas on leaves around feeding site

Identification

- Adults: Larger than our native stink bugs, this bug can be 12 - 17 mm long and has a base color of brown, red, and black on the dorsal surface. A distinguishing factor of adults are white bands on the legs and antennae and alternating dark and light bands on the abdomen.
- Nymphs: Hatching at 2.4 mm in length with black heads and orange abdomens, they transition from orange red to brownish and begin to develop the white bands on legs and antennae and spines on the humeral margins.

Image Credits: Top Left: Mike Lewis, Center for Invasive Species Research, UC Riverside; Top Right: Bryce Sutherland, S&R Ag Consulting; Bottom Left: Hectonichus, Wikipedia; Bottom Right: Gary Bernon, USDA APHIS



Adult on fruit



Fruit Damage

5560937



Fifth to sixth instar nymph



Newly hatched eggs and nymphs



Spongy Moth (*Lymantria dispar*)



NOT SHOWN TO SCALE



Brown Male Moth and White Female Moth, Image Credit: John Ghent

Hosts (this list is not exhaustive)

Preferred hosts: Oaks (*Quercus spp.*)
Apple, alder, basswood, birch, larch, poplar, sweet gum, willow, hawthorn

Signs and Symptoms

- Defoliated leaves
- Heavily infested trees will look barren in the summer
- Egg masses on bark or can be found on outdoor object
- Tan masses of eggs with hairy covering being the size of a quarter to 7.6 cm long

Identification

- Adults: Males have a 3.8 cm wingspan and are brown with dark brown patterns on the wings. Females are nearly white with zig-zag patterns on their wings, 6.4 cm wingspan, but are flightless. Moths can be seen in the summer in July and August.
- Larvae: Caterpillars are 3.8 - 6.4 cm long. Newly hatched, caterpillars start black and hairy and molt to gray as they mature with bristled tufts of hair and a pattern with 5 pairs of blue dots then 6 pairs of red dots along their backs. Caterpillars can be found in the spring around April- June.

Image Credits: Top Left: Ferenc Lakatos, University of Sopron; Top Right: Karla Salp, Washington State Department of Agriculture, Bugwood.org; Bottom Left: Steven Katovich, Bugwood.org; Bottom Right: John Ghent, Bugwood.org



Pupae



Adult Female on Egg Mass

5551083



Caterpillar



Adult Male



Elongate Hemlock Scale (*Fiorinia externa*)



Infestation. Image Credit: Eric R. Day, Virginia Polytechnic Institute and State University

Hosts

Hemlocks (*Tsuga spp.*) and firs (*Abies spp.*)
Occasionally: Spruce (*Picea spp.*), pines (*Pinus spp.*),
and Douglas fir (*Pseudotsuga menziesii*)

Signs and Symptoms

- Yellow spotting on needles
- Premature needle loss
- Branch and limb dieback
- Thin crowns (More typical with severe infestations)
- Scales covering the underside of the needles

Identification

- Adults: These insects are barely visible to the naked eye and may go unnoticed.
- Females are soft-bodied, legless, and wingless and are only 2 mm long. The yellow, immobile females lay their eggs under an orange, brown wax coating.
- Males are light brown, 1.5 mm long and may look like tiny wasp parasitoids and can fly. Because they have two overlapping generations a year, all life stages can be found year-round.

Image Credits: Top Left: Eric R. Day, Virginia Polytechnic Institute and State University; Top Right: John A. Davidson, University of Maryland College Park; Bottom Left: Kristen Wickert, USDA Forest Service; Bottom Right: Kristopher Abell, University of Massachusetts



Infestation



White male cover and brown female covers.



Sign(s) of Infestation



Adult(s)



Phytophthora species



Symptomatic leaf affected by *P. ramorum*. Photo Credit: Joseph OBrien, USDA Forest Service

Signs and Symptoms

- Small, chlorotic, or wilting leaves
- Leaf blight
- Thin and sparse crowns
- Branch, twig, or shoot dieback
- Heavy cone production
- Cankers
- Trees may have bleeding lesions on the trunk or from cankers
- The inner bark beneath the lesions can be orange to pink to brown in color

- Root rot
- Decreased fruit size and yield
- Necrotic lesions that start as blackening of leaf petiole and can extend to the base of the leaf, potentially progressing to cover the entire leaf

Spread

Phytophthora spp. can be spread through human movement of contaminated plant materials, soil, or tools. *Phytophthora* is able to spread through water, allowing it to move within soil and infect uninfected plants.

Image Credits: Top Left: Forest Research, United Kingdom Forestry Commission, Bugwood.org; Top Right: Joseph Hulbert, WSU; Bottom Left: Joseph OBrien, USDA Forest Service, Bugwood.org; Bottom Right: Sarah Green, Forest Research



Bleeding on bark surface of coast live oak, caused by *P. ramorum*



Stem lesions caused by *P. ramorum* on Tanoak



Canker caused by *P. cinnamomi* on oak



P. austrocedri causing juniper tree dieback



Phytophthora ramorum

Affects native oak and tanoak forests in Southern Oregon and California and causes lethal stem cankers, also known as sudden oak death. Besides infecting oak, *P. ramorum* is known to have a wide range of hosts and can spread aerially.


Forest trees killed by P. ramorum in Oregon. Photo Credit: Joseph Hulbert, WSU.



Phytophthora cinnamomi

Known to have the broadest host range among *Phytophthora* species. It has a global distribution, causing extensive damage in forests in Australia, the Mediterranean, and southeast United States. It tends to cause the largest issues in warmer climates.

Western Australia landscape decimated by P. cinnamomi. Photo Credit: Trudy Paap, FABI.



Phytophthora occultans

Causes root, collar, and crown rot, and stem cankers on woody plants in multiple plant families. *P. occultans* was originally found in the Netherlands and has since been found in other parts of Europe and the United States.

Dead sword fern in container where P. occultans was found in Tacoma, Washington. Photo Credit: Joseph Hulbert, WSU



Phytophthora austrocedri

The cause of “mal del ciprés” the disease affecting *Austrocedrus chilensis* in Andes mountain range of Chile and southern Argentina. Also known to affect common juniper (*Juniperus communis*) and Mediterranean cypress (*Cupressus sempervirens*). It was recently found in an Oregon nursery.

Dead and dying Austrocedrus chilensis (Cordilleran cypress) affected by P. austrocedri. Photo credit: E.M. Hansen, OSU



Phytophthora alni

Phytophthora alni is a concern for alder (*Alnus* spp.) populations because it causes lethal root and collar rot. The disease has been found across several countries in Europe, but it is highly variable likely due to *P. alni* hybridizations with other species. It is a USDA APHIS Priority Pest.

Disease symptoms seen under bark of alder trees (alnus spp.). Photo credit: Andrej Kunca, National Forest Centre Slovakia



Phytophthora kernoviae

Phytophthora kernoviae can affect many plant species. It is well known in the United Kingdom because it causes lesions on rhododendron leaves and European beech stems. It is also a USDA APHIS Priority Pest.

Wilting rhododendron leaves affected by P. kernoviae. Photo credit: Forest Research, United Kingdom Forestry Commission



Laurel Wilt (*Rafaelea lauricola*)



Felling dead redbays. Image Credit: Albert Mayfield, USDA Forest Service

Hosts

Oaks (*Quercus spp.*), *lauracea*

Causal Agent

Rafaelea lauricola fungus

Signs and Symptoms

- Tree wilt starts in one portion of a tree and eventually spreads
- Defoliation or browning/ reddening of leave depending on the host
- Black staining of the sapwood

Spread

Laurel wilt is caused by the fungal pathogen, *Rafaelea lauricola*, spread through the Redbay ambrosia beetle (*Xyleborus glabratus*). As the fungus spreads, the water-conducting cells become clogged.

Redbay Ambrosia Beetle

Identification

- Cylinder shaped
- Brown to black color
- 2 mm long
- 1 mm diameter entrance holes

Image Credits: Top Left: Chip Bates, Georgia Forestry Commission; Top Right: Albert (Bud) Mayfield, USDA Forest Service, Bugwood.org; Bottom Left: Mary Ann Hansen, Virginia Polytechnic Institute and State University; Bottom Right: Joseph Benzel, Screening Aids, USDA APHIS



Infection on camphor tree



Redbay with black staining of sapwood



Redbay with black staining of sapwood



Redbay Ambrosia Beetle



Ash Dieback (*Hymenoscyphus fraxineus*)



Ash dieback , Image Credit: Forest Research, United Kingdom

Hosts

Ash (*Fraxinus spp.*)

Causal Agent

Hymenoscyphus fraxineus fungus

Spread

Hymenoscyphus fraxineus is spread locally by wind, but the movement of diseased ash plants is causing more widespread dispersal of this pathogen.

Signs and Symptoms

- Leaf wilting and discoloration
- Early leaf drop
- Lesions where branches meet the trunk
 - Diamond shaped
- Inner bark under lesions is brown to gray
- Epicormic growth

Image Credits: Top Left: Cumbria Wildlife Trust; Top Right: Ludwig Treuter, iNaturalist; Bottom Left: Joe Bates, Woodland Trust; Bottom Right: Thomas Kirisits, Forest Research United Kingdom



Black lesion on ash



Hymenoscyphus fraxineus on dead leaf stalk



Ash leaf wilting



Small, dried lesion



Signs of oak wilt. Image Credit: Ryan Armbrust, Kansas Forest Service, Bugwood.org

Hosts

Oak (*Quercus spp.*)

Causal Agent

Bretziella fagacearum fungus

Spread

Oak wilt can spread locally through connected root systems. The fungus can also be transferred from an infected to a healthy tree through oak bark beetles (*Scolytidae*) and sap beetles (*Nitidulidae*) which can carry spores to tree wounds. Human movement is another vector that can spread oak wilt.

Signs and Symptoms

- Flagging in upper canopy
- Leaves become dull green and wilted
- Bronze or brown tissue at leaf tip and margins
- Defoliation
- Black/gray fungal mats under the desiccating bark
- Sapwood streaking

Image Credits: Top Left & Top Right: Steven Katovich, Bugwood.org; Bottom Left: alexfeltmeyer1, iNaturalist; Bottom Right: Sandra Jensen, Cornell University, Bugwood.org



Large yard tree killed by oak wilt



Oak wilt pocket



Bretziella fagacearum on Northern red oak



Dark streaking on wood of oak branch



Sooty Bark Disease on big leaf maple, Image Credit: Joey Hulbert, WSU

Hosts

Maples (*Acer spp.*)

Horse chestnut (*Aesculus hippocastanum*)

Causal Agent

Cryptostroma corticale fungus

Spread

Sooty bark disease spores can spread through wind and rain. Humans can aid in the movement of the fungus through diseased plant materials, but it is hypothesized to be native and widespread throughout the northwest.

Signs and Symptoms

- Crown dieback
- Small or dwarfed leaves
- Black 'sooty' fungal crust within the bark (not on the xylem tissues or outer bark)
- White spores within the bark when the outer bark is pushed off
- Staining within the xylem tissues

Image Credits: Top Left & Top Right: Faith Dewitz, WSU, Bottom Left: & Bottom Right: Joey Hulbert, WSU



Old patch of sooty bark with new, white spores



Spores underneath big leaf maple bark



Staining inside the infected sycamore maple



'Sooty' patch on big leaf maple



American Elm with symptoms of Dutch Elm Disease. Image Credit: Ward Upham, Kansas State University

Hosts

Elm (*Ulmus spp.*)

Causal Agent

Ophiostoma ulmi and *Ophiostoma novo-ulmi* fungi

Spread

Dutch elm disease spreads through root grafts and elm bark beetles, including beetles from the *Scolytus* genus and the native elm bark beetle (*Hylurgopinus rufipes*)

Signs and Symptoms

- Leaves or branches in the outer crown yellow, wilt, then turn brown
- Discoloration and wilting progresses towards the tree trunk
- Premature leaf drop in spring and summer
- Diseased foliage and shoots
- Shoot tip die back
- Brown streaks in sap wood

Image Credits: Top Left: Steven Katovich, Bugwood.org; Top Right: William Jacobi, Colorado State University; Bottom Left: M. Grabowski, UMN Extension; Bottom Right: Roland J. Stipes, Virginia Polytechnic Institute and State University, Bugwood.org



Multiple egg galleries of native elm bark beetle



Brown staining of sapwood



Flagging of leaves on an infected tree



Wilted American elm leaves



Beech Leaf Disease (*Litylenchus crenatae*)



Beech leaf disease on American Beech , Image Credit: Matthew Borden, Bartlett Tree Experts, Bugwood.org

Hosts

Beech (*Fagus spp.*)

Causal Agent

Litylenchus crenatae subsp. *mccannii* nematode

Spread

Nematodes (microscopic roundworm) that cause beech leaf disease are primarily spread through human movement of infected plant materials.

Signs and Symptoms

- Dark bands on leaves in early spring caused by nematodes
- Leathery, deformed, crinkled leaves
- Yellowing leaves as season progresses
- Reduced bud and leaf production as the disease progresses
- Premature leaf drop
- Canopy dieback and thinning

Image Credits: Top Left: Paulo Vieira, USDA, ARS; Top Right & Bottom Left: Matthew Borden, Bartlett Tree Experts; Bottom Right: Yonghao Li, The Connecticut Agricultural Experiment Station, Bugwood.org



Litylenchus crenatae *mccannii* nematode under microscope



Damage on new leaves



Symptoms and damage on beech leaves



Banding symptoms on leaf

REPORT SIGHTINGS

Share reports with the QR codes
websites, or contacts below.

Washington



invasivespecies.wa.gov

Oregon



oregoninvasiveshotline.org

MORE RESOURCES

Forest Health Watch

Participate as community scientists, share reports,
or contact the program staff at:

<https://foresthealth.org>

WSU Urban Forest Health Lab

Find more information, trainings and resources at

<https://treehealth.wsu.edu>

