



Evaluation of Changes in Vegetation Indices Due to In- field Sensor Angles

Maicynn Hansen


University of Illinois at Urbana-Champaign

Phenomix Big Data Management


Mentor: Dr. Sindhuja Sankaran

Strategies to Improve Production

- Fertility
- Pest/Disease Control
- Precision/Other Management
- **Crop Breeding**
 - Phenotyping is often demanding and time intensive



<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0066428>





Introduction

- Phenomics is useful for large scale phenotyping
- RGB and multispectral imagery capture crop reflectance measurements
- Vegetation indices (VIs) based on reflectance data provides insight into crop responses



Internet of Things (IoT) Sensor System




Worasit Sangjan



Introduction

- IoT sensor systems enable real time imaging and management decisions



Does the sensor angle affect the estimated crop vegetation indices?

Methods

- Crops: Winter wheat & spring pea
- IoT data angles: 30°, 45°, 60°
- Drone data: at nadir (ground-reference data)
- Data type: RGB and NoIR images




Image acquisition

Image separation




Radiometric correction




Plot segmentation

<https://www.oreilly.com/library/view/hands-on-image-processing/>



Histogram analysis




Methods

Multispectral

- Green normalized difference vegetation index (GNDVI)


RGB

- Normalized green red difference index (NGRDI)
- Green leaf index (GLI)



Extraction of vegetation indices

Statistical analysis




- ANOVA test for comparison between angles
- Pearson correlation for comparison between drone with IoT data


Spring Pea Results

The angles of 30° and 45° are most similar for GLI and NGRDI

ANOVA, $p = 2.2e-16$ ***




Drone ground reference data had the strongest correlation with images from 45°




Winter Wheat Results

30° and 45° were most similar for all three vegetation indices



Drone ground reference data had the strongest correlation with 60°



Discussion

- Angles have an affect on calculated vegetation indices
- Image processing methods can be better optimized for data extraction


Challenges

- Soil
- Plant senescence
- Reference panel position



Future Work

Distortion correction techniques to correct inaccurate VIs



Acknowledgements

Worasi Sangjan

Milton Valencia Ortiz

Dr. Arron H. Carter

Dr. Rebecca J. McGee



Summer 2021

**FACT: Research Experience for Undergraduates on Phenomics Big
Data Management**



**United States Department of Agriculture
National Institute of Food and Agriculture**

**Research and Extension Experiences for Undergraduates (REEU)
Project: 1021788**



Questions?

