

Calculations of Electronic Stopping Cross Sections for Low-energy Protons in Water

S. Uehara¹, L. H. Toburen², W. E. Wilson³, D. T. Goodhead⁴, H. Nikjoo⁴

¹School of Health Sciences, Kyushu University, Maidashi 3-1-1, Higashi-ku, Fukuoka 812-8582, Japan; ²Department of Physics, East Carolina University, Greenville, ND 27858, USA; ³US Transuranic and Uranium Registries, Washington State University, Tri-Cities, Richland, WA 99352, USA; ⁴Medical Research Council (MRC) Radiation and Genome Stability Unit, Harwell, Oxfordshire OX11 0RD, UK

Calculations of electronic stopping cross sections for low-energy protons were performed using available interaction cross-sections and information on the mean energy loss per collision. The results were compared with published stopping power data to assess the general consistency and reliability of the results. The calculated stopping cross sections include contributions from excitation by both the proton and neutral-hydrogen components of the charge-equilibrated beam and the mean energy lost per interaction. The calculated electronic stopping cross sections for swift charge-equilibrated hydrogen particles in water are in agreement with previous tabulations of stopping power to within 20% (approx.) for particles energies less than 0.3 MeV.

Published in Radiation Physics and Chemistry Vol. 59: 1-11; (2000)

USTUR-0130-98