

59th Annual Meeting of the Health Physics Society
Baltimore, Maryland
July 13 – 17, 2014

USTUR Whole-Body Case 0212: Testing NCRP Wound Model

Maia Avtandilashvili, Stacey L. McComish
and Sergei Y. Tolmachev

US Transuranium and Uranium Registries
College of Pharmacy, Washington State University
Richland, WA 99354



*“Learning from Plutonium
and Uranium Workers”*

WASHINGTON STATE
UNIVERSITY
World Class. Face to Face.

Motivation

- ❑ >2,000 contaminated wounds reported
 - Commonly treated by tissue excision and chelation
- ❑ NCRP 156 Wound Model (2007)
 - Based exclusively on animal experiments
 - Important to test against human data
- ❑ Effectiveness of chelation treatment
 - Residual vs. projected dose
- ❑ USTUR Whole Body donations: *A total of 42 cases*
 - 10 cases with documented wound intakes: ^{239}Pu (9), ^{241}Am (1)
 - 4 cases with a single Pu wound as a major intake
 - Case 0212 selected for this study




USTUR Whole Body Case 0212

- Exposure: Wound (treated)
- Treatment: Tissue Excision & Ca-DTPA
- Donation Year: 1984
- Post-Intake: 17 y
- Cause of Death: Pulmonary Emphysema
- Age: 56 y




Contaminated Wound

- ❑ Left middle finger, anterior surface
- ❑ Cutting a pipe on the solvent extraction hood
- ❑ Material involved: $\text{Pu}(\text{NO}_3)_4$
- ❑ Initial survey meter reading: 10,000 dpm
 - Decontaminated to 500 dpm
- ❑ Initial wound count: 59 nCi (~2.2 kBq)
 - After wound excision: 11 nCi (~0.4 kBq)
 - Excised tissue count: 122 nCi (~4.5 kBq)
- ❑ Chelation treatment: 6 months – bi-weekly
 - A total of 26.5 g Ca-DTPA




Urinalysis Data

- Analysis Method:
 - ✓ Autoradiography
- MDA: $\sim 0.001 \text{ Bq d}^{-1}$
- A total of 205 samples




Urinalysis Data: 'Treated'

- Analysis Method:
 - ✓ Autoradiography
- MDA: $\sim 0.001 \text{ Bq d}^{-1}$
- A total of 205 samples
- 180 samples
 - ✓ affected by DTPA
 - ✓ 201 days post-intake
- $\sim 916 \text{ Bq Pu}$ excreted during treatment
 - ✓ Max rate: 73 Bq d^{-1}



Urinalysis Data: 'Untreated'

- Analysis Method:
 - ✓ Autoradiography
- MDA: $\sim 0.001 \text{ Bq d}^{-1}$
- A total of 205 samples
- 25 samples
 - ✓ not affected by DTPA
- Ave. post-DTPA rate:
 - ✓ $0.02 \pm 0.01 \text{ Bq d}^{-1}$



Autopsy Tissue Sample Analysis


- Total of 264 tissue samples
- Analysis Method: *Alpha Spectrometry*

Tissue	Concentration, Bq kg ⁻¹	Activity, Bq
Wound (<i>muscle & skin</i>)	40.2 ± 0.7	14.3 ± 0.3
Lungs including LNTH	0.48 ± 0.07	0.82 ± 0.12
Skeleton (<i>174 samples</i>)	11.0 ± 0.1	114.5 ± 0.5
Liver	33.8 ± 1.1	80.5 ± 2.6
Kidneys	0.37 ± 0.01	0.172 ± 0.005
Soft Tissues (<i>56 samples</i>)	0.0004 ± 0.0001	34.2 ± 0.6
Total Systemic	n/a	229.2 ± 2.7




Methods

- ❑ Internal Dosimetry Software
 - IMBA Professional Plus[®]
- ❑ Maximum likelihood fitting of:
 - ‘Baseline’ (post-treatment) urine data
- ❑ Models applied:
 - ICRP 67 Pu Systemic Model
 - NCRP Wound Model





NCRP Wound Model



- ☐ Default material types:
- Soluble
 - ✓ Weak
 - ✓ Moderate
 - ✓ Strong ← $\text{Pu}(\text{NO}_3)_4$
 - ✓ Avid
 - Colloid
 - Particle
 - Fragment



Results: Excretion




☐ Credible fit:

➤ χ^2 alpha = 0.246



Results: Retention



Pu Retention at Time of Death, Bq

Organ	Predicted	Measured
Liver	97.3	80.5
Skeleton	147.4	114.5
Wound	6.8	14.3

Model predicted:

- ❑ Pu in liver and skeleton
 - within ~25%
- ❑ Pu in wound
 - by a factor of 2



Results: Intake and Dose

□ Intake


➤ IMBA estimate:

364 Bq

➤ Pu excreted with DTPA:

916 Bq

1,280 Bq



□ Committed Effective Dose

➤ Residual CED: 177 mSv

vs.

➤ Projected CED: 622 mSv



Future Work

- ❑ Apply Bayesian analysis methods:
 - Define suitable priors for wound model parameters
 - Use Monte Carlo simulation (**WeLMoS**, **MCMC**) to:
 - ✓ derive best estimate of intake
 - ✓ calculate uncertainties in wound retention parameters
 - ✓ quantify effectiveness of DTPA treatment



Questions?

