

Internal Dosimetry Session  
62<sup>nd</sup> Annual Meeting of the Health Physics Society  
Raleigh, NC, July 9 - 13, 2017

# Enhancement of Plutonium Excretion Following Late Ca-EDTA/DTPA Treatment

Sara Dumit, Maia Avtandilashvili, Sergei Y.  
Tolmachev

United States Transuranium and Uranium Registries  
College of Pharmacy, Washington State University

*“Learning from Plutonium and  
Uranium Workers”*






# Objectives


- Calculate plutonium excretion enhancement following delayed Ca-EDTA/DTPA treatment
- Compare effectiveness of chelating agents
- Evaluate Pu-EDTA complex removal half-time

# Chelation Treatment for Plutonium

- Chelating agents
- Remove radioactive elements from the body
- Enhance metal excretion by forming stable complex
- Chelating agents for Pu decorporation:



Ca-EDTA  
ethylenediaminetetraacetic  
acid



Ca-DTPA  
diethylenetriaminepentaacetic  
acid



# Chelation Treatment: Dosimetry Challenge

- Administration of chelating agents alters radionuclide biokinetics
- Urinary excretion of radionuclide is enhanced significantly
- Common practice for radiation dose estimation is to exclude data affected by treatment
- Affected data may be corrected using enhancement factor



# Enhancement Factor

- Enhancement factor (EF)

*measured radionuclide daily excretion rate at the day of chelation divided by expected excretion on the same day without chelation (LaBone 1994)*

- Empirical parameter determined on case-by-case basis. Varies significantly among individuals
- Recommended value for plutonium decorporation with Ca-DTPA is 50



# USTUR Case 0785 Summary

- Primary exposure: Plutonium (Pu)
- Exposure scenario: Acute inhalation and wound
- Material type: Mixture (M and S)
- Treatment: Wound excision  
Ca-EDTA/DTPA chelation
- Smoking status: Yes (cigars)
- Cause of death: Mesothelioma
- Post-intake: 51 y
- Age: 79 y




# Ca-EDTA/DTPA Treatment Regimen

- (I) Initial: Ca-EDTA, 67g (daily)
- Week 1:  $1\text{g} \times 2 \times 7\text{d}$
  - Week 2:  $1\text{g} \times 2 \times 3\text{d}; 2\text{g}+1\text{g}; 2\text{g} \times 2 \times 1\text{d}$
  - Weeks 3 – 4:  $2\text{g} \times 2 \times 10\text{d}$
- (II) Delayed: Ca-EDTA, 40g (daily)
- Week 39 – 40:  $2\text{g} \times 2 \times 10\text{d}$
- (III) Delayed: Ca-DTPA, 12g (weekly)
- Weeks 374 – 385:  $1\text{g} \times 1 \times 12\text{wk}$



# Plutonium Urinary Excretion






# Ca-EDTA Enhancement Factor

Regimen II: weeks 39 - 40

- N = 10
- Range: 38 - 277
- Mean:  $150 \pm 70$






# Ca-DTPA Enhancement Factor

Regimen III: weeks 374 - 385

- N = 11
- Range: 25 - 134
- Mean:  $74 \pm 35$





# Effectiveness of Chelating Agent

- The measure of effectiveness,  $E_{\text{DRUG}}$


$$E_{\text{DRUG}} = \frac{A}{D \times t}$$

$A$  (Bq) - activity excreted during time  $t$

$D$  (g) - dosage of an administered drug



$t$  (d) - duration of the treatment plus 100 days

# Effectiveness: Ca-EDTA vs Ca-DTPA





# Pu-EDTA Complex Removal





# Summary

- Plutonium excretion enhancement factors were calculated following delayed Ca-EDTA/DTPA treatment. For Ca-EDTA, values ranged from 38 - 277 with a mean of  $150 \pm 70$ ; for Ca-DTPA, values ranged from 25 - 134 with a mean of  $74 \pm 35$
- Ca-EDTA enhancement factors for 5 daily injections showed monotonic increase from day 1 to 4, followed by decrease on day 5
- Attempt was made to compare effectiveness of chelating agents. Delayed administration of Ca-EDTA was significantly less effective than initial; delayed administration of Ca-DTPA was as effective as initial Ca-EDTA
- The half-time of Pu-EDTA complex removal in urine was evaluated to be 1.6 d which was comparable with 3.2 d estimated for USTUR Case 0269



Thank you!

Questions?