IDEQ Machine Learning Air Quality Forecast System Update

NW-AIRQUEST Meeting

02/03/2022

Overview of Forecast System

Spatial Coverage

O3 Sites (50)



PM2.5 Sites (130)



Site Selection Criteria: Having at least 3 years of data and continue to operate in the future

Overview of Forecast System

Data Process – In and Out



UW Meteorology Models

- Currently 26 ensemble members
- Initialized at 00Z and 12Z
- Each forecasts 72 hours
- Resolution: 4km

- Time of arriving
  - 00Z forecast : ~09 - 12 MST
  - 12Z forecast : ~21 - 00 MST
- Utilization
  - Morning forecast : using previous day 12Z Met forecast
  - Afternoon forecast : using current day 00Z Met forecast

O3 and PM2.5 Forecast
Machine Learning Models Used

Neural Network Models

- Dense Neural Network Model
- 1D Convolutional Neural Network Model
- Recurrent Neural Network (LSTM)

Decision Tree Based Models

- Gradient Boosting
- Random Forest
- Boosted Random Forest

Baseline Model : Persistence

O3 and PM2.5 Forecast Inputs Used

11 Inputs :

- Previous 24th hour O3/PM2.5
- T, P, RH, PBLH, WS, WDIR
- O3/PM2.5 hourly_mean (grouped by Month, Weekday/end, Hour)
- Month (1-12), Weekday (0-1), Hour (0-23)

Ensemble

- Three layers of Ensemble
- Ensemble already employed in some Machine Learning Models, such as random forest
- Ensemble of Machine Learning Models:
  - Produce the final model output from multiple machine learning models for one set of input data
- Ensemble of Meteorology Models
  - The distribution of prediction

Bias Correction for Day 1

- Replace prediction with known observation
- Apply bias correction to directly following several hours based on known bias of previous 3 hours
- New treatment : Apply bias correction for the rest hours based on previous day's 3-hour window hourly bias



Retraining Model Annually

- Why
  - Machine Learning Model needs a lot of data
  - Model Performance may drift overtime
- When and How
  - Each January
  - Using previous year's data as test data set and free up part of original test data into training data set
  - Tune up model parameters

Model Performance
at
St. Lukes Meridian
for
O3 Models Used in Year 2021

St. Lukes Meridian O3 Site
Year 2021 Model Performance (1)

Hourly Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	52	8.99	144.53	0	0.3	0.46	12.02
pure_model_am_day2	48	7.19	81.76	-0.1	0.24	0.69	9.04
pure_model_pm_day2	47	7.15	81.13	-0.1	0.24	0.69	9.01
pure_model_am_day1	47	6.98	77.18	-0.1	0.23	0.71	8.79
pure_model_pm_day1	46	7.1	79.08	-0.12	0.24	0.7	8.89
bias_corrected_am_day1	47	4.66	48.54	-0.07	0.16	0.82	6.97
bias_corrected_pm_day1	45	2.95	31.19	-0.03	0.1	0.88	5.59

Daily Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	37	5.72	62.67	0	0.13	0.66	7.92
pure_model_am_day2	23	5.98	57.03	-0.09	0.14	0.68	7.55
pure_model_pm_day2	22	5.76	53.18	-0.08	0.14	0.71	7.29
pure_model_am_day1	22	5.61	50.8	-0.08	0.13	0.72	7.13
pure_model_pm_day1	22	5.69	51.55	-0.09	0.13	0.72	7.18
bias_corrected_am_day1	22	4.86	40.27	-0.06	0.11	0.78	6.35
bias_corrected_pm_day1	17	3.03	17.71	-0.02	0.07	0.9	4.21

St. Lukes Meridian O3 Site
Year 2021 Model Performance (2)

AQI Classification Metrics for All AQI Class

Forecast	AQI_class	accuracy	HSS	KSS
persistence	ALL	0.86	0.5	0.5
pure_model_am_day2	ALL	0.9	0.58	0.51
pure_model_pm_day2	ALL	0.89	0.54	0.47
pure_model_am_day1	ALL	0.9	0.58	0.52
pure_model_pm_day1	ALL	0.89	0.55	0.49
bias_corrected_am_day1	ALL	0.9	0.58	0.52
bias_corrected_pm_day1	ALL	0.89	0.59	0.61

AQI Classification Metrics for AQI Class 2 (Yellow)

Forecast	AQI_class	precision	recall	f1-score	support
persistence	AQI class 2	0.5	0.5	0.5	48
pure_model_am_day2	AQI class 2	0.66	0.58	0.62	43
pure_model_pm_day2	AQI class 2	0.61	0.53	0.57	47
pure_model_am_day1	AQI class 2	0.65	0.58	0.62	48
pure_model_pm_day1	AQI class 2	0.63	0.54	0.58	48
bias_corrected_am_day1	AQI class 2	0.65	0.58	0.62	48
bias_corrected_pm_day1	AQI class 2	0.57	0.69	0.62	48

St. Lukes Meridian O3 Site
Year 2021 Hourly Time Series





Three “normal” days followed by wildfire impact days”

St. Lukes Meridian O3 Site
Year 2021 Daily Scatter Plots



Model Performance
at
St. Lukes Meridian
for
Retrained O3 Models

St. Lukes Meridian O3 Site
Year 2021 Model Performance
Original vs. Retrained(1)

Original Model Hourly Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	52	8.99	144.53	0	0.3	0.46	12.02
pure_model_am_day2	48	7.19	81.76	-0.1	0.24	0.69	9.04
pure_model_pm_day2	47	7.15	81.13	-0.1	0.24	0.69	9.01
pure_model_am_day1	47	6.98	77.18	-0.1	0.23	0.71	8.79
pure_model_pm_day1	46	7.1	79.08	-0.12	0.24	0.7	8.89
bias_corrected_am_day1	47	4.66	48.54	-0.07	0.16	0.82	6.97
bias_corrected_pm_day1	45	2.95	31.19	-0.03	0.1	0.88	5.59

Retrained Model Hourly Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	52	9	144.76	0	0.3	0.45	12.03
pure_model_am_day2	44	6.9	76.27	-0.1	0.23	0.71	8.73
pure_model_pm_day2	43	6.78	74.26	-0.1	0.23	0.72	8.62
pure_model_am_day1	43	6.68	72.04	-0.1	0.22	0.73	8.49
pure_model_pm_day1	42	6.74	72.71	-0.1	0.23	0.73	8.53
bias_corrected_am_day1	37	4.3	46.32	-0.04	0.14	0.83	6.81
bias_corrected_pm_day1	40	2.67	28.19	-0.02	0.09	0.89	5.31

St. Lukes Meridian O3 Site
Year 2021 Model Performance
Original vs. Retrained(2)

Original Model Daily Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	37	5.72	62.67	0	0.13	0.66	7.92
pure_model_am_day2	23	5.98	57.03	-0.09	0.14	0.68	7.55
pure_model_pm_day2	22	5.76	53.18	-0.08	0.14	0.71	7.29
pure_model_am_day1	22	5.61	50.8	-0.08	0.13	0.72	7.13
pure_model_pm_day1	22	5.69	51.55	-0.09	0.13	0.72	7.18
bias_corrected_am_day1	22	4.86	40.27	-0.06	0.11	0.78	6.35
bias_corrected_pm_day1	17	3.03	17.71	-0.02	0.07	0.9	4.21

Retrained Model Daily Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	37	5.71	62.53	0	0.13	0.66	7.91
pure_model_am_day2	21	5.57	49.92	-0.08	0.13	0.72	7.07
pure_model_pm_day2	21	5.33	45.93	-0.07	0.13	0.75	6.78
pure_model_am_day1	25	5.23	45.86	-0.07	0.12	0.75	6.77
pure_model_pm_day1	23	5.26	45.75	-0.08	0.12	0.75	6.76
bias_corrected_am_day1	28	4.23	34.16	-0.04	0.1	0.81	5.84
bias_corrected_pm_day1	14	2.6	12.96	-0.01	0.06	0.93	3.6

St. Lukes Meridian O3 Site
Year 2021 Hourly Time Series
Original vs. Retrained

Original



Retrained



Retrained Model for St. Lukes Meridian O3 Site Year 2021 Daily Scatter Plots

Original



Retrained



Model Performance
at
St. Lukes Meridian
for
PM2.5 Models

St. Lukes Meridian PM2.5 Site
Year 2021 Model Performance (1)

Hourly Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	137	5.84	94.7	0.01	0.61	0.24	9.73
pure_model_am_day2	134.9	4.8	70.93	-0.04	0.5	0.43	8.42
pure_model_pm_day2	135.6	4.68	67.02	-0.03	0.49	0.46	8.19
pure_model_am_day1	135.5	4.6	64.4	-0.04	0.48	0.48	8.03
pure_model_pm_day1	135.1	4.61	63.85	-0.03	0.48	0.49	7.99
bias_corrected_am_day1	106	3.1	41.27	-0.02	0.32	0.67	6.42
bias_corrected_pm_day1	106.1	2	25.42	0.01	0.21	0.8	5.04

Daily Regression Performance Metrics

Forecast	max_error	mean_absolute_error	mean_squared_error	normalized_mean_bias	normalized_mean_error	r2_score	root_mean_squared_error
persistence	36.1	3.6	35.23	0.01	0.38	0.62	5.94
pure_model_am_day2	57.6	3.48	40.71	-0.04	0.36	0.56	6.38
pure_model_pm_day2	52.7	3.25	36.86	-0.03	0.34	0.6	6.07
pure_model_am_day1	49.8	2.99	30.75	-0.04	0.31	0.67	5.55
pure_model_pm_day1	49	3.04	30.62	-0.03	0.32	0.67	5.53
bias_corrected_am_day1	25.9	2.02	12.63	-0.02	0.21	0.86	3.55
bias_corrected_pm_day1	21.1	1.49	6.83	0.01	0.16	0.93	2.61

St. Lukes Meridian PM2.5 Site Year 2021 Model Performance (2)

AQI Classification Metrics for All AQI Class

Forecast	AQI_class	accuracy	HSS	KSS
persistence	ALL	0.83	0.5	0.51
pure_model_am_day2	ALL	0.83	0.5	0.5
pure_model_pm_day2	ALL	0.85	0.56	0.56
pure_model_am_day1	ALL	0.84	0.53	0.52
pure_model_pm_day1	ALL	0.84	0.54	0.54
bias_corrected_am_day1	ALL	0.91	0.73	0.73
bias_corrected_pm_day1	ALL	0.92	0.78	0.78

AQI Classification Metrics for AQI Class 2 (Yellow)

Forecast	AQI_class	precision	recall	f1-score	support
persistence	AQI class 2	0.54	0.56	0.55	66
pure_model_am_day2	AQI class 2	0.55	0.57	0.56	65
pure_model_pm_day2	AQI class 2	0.6	0.63	0.62	65
pure_model_am_day1	AQI class 2	0.6	0.59	0.6	66
pure_model_pm_day1	AQI class 2	0.59	0.62	0.6	66
bias_corrected_am_day1	AQI class 2	0.76	0.76	0.76	66
bias_corrected_pm_day1	AQI class 2	0.83	0.79	0.81	66

St. Lukes Meridian PM2.5 Site Year 2021 Time Series (1)



St. Lukes Meridian PM2.5 Site
Year 2021 Time Series (2)



St. Lukes Meridian PM2.5 Site
Year 2021 Time Series (3)



St. Lukes Meridian PM2.5 Site
Year 2021 Daily Scatter Plots



St. Lukes Meridian PM2.5 Site A Peek of Model Performance in Year 2022





Overview of Forecast System

Data Process – In and Out



Demo

- Email Notification
- Simple Online Map
  - Make it easy to navigate the sites
- Spreadsheet for each Regional Office
  - Regional offices pick the sites

Path Forward

- Bring in NASA GEOS-FP model output as new inputs to improve our model performance in wildfire and dust storm impact period
- Evaluate the model performance in January of each year
- Retrain the machine learning models with last year's data added in January of each year
- Test and bring in new inputs in the future

Questions and Discussion

The End

Supplemental Slides

Email Notification

Machine Learning Air Quality Forecast is here! (Forecasted at 01/26/2022 Morning)

WZ Wei.Zhang@deq.idaho.gov
To Wei Zhang

Reply | Reply All | Forward | ...
Wed 1/26/2022 7:46 AM

Wei,

This morning's forecast is here! The result can be found at:

http://10.220.98.54/ml_forecast_outputs/2022/20220126_am/

OR through simple online map http://10.220.98.54/ml_forecast_outputs/2022/20220126_am/20220126_am_forecast_0Map.html

To access forecasts for previous days, please go to:

http://10.220.98.54/ml_forecast_outputs/

This forecast is based on a machine learning method utilizing modeled ensemble meteorological forecasts from the University of Washington. Acute or event-based impacts, such as wildfires and dust storms, are not considered in the model. This forecast should be used as a starting point and then adjusted based on local knowledge of these factors.

Hopefully this tool makes your life easier. Please provide us any feedback you may have.

Technical Services Division
Idaho Department of Environmental Quality
Wei.Zhang@deq.idaho.gov

Simple Online Map




PM2.5 Daily 24 Hour Average Concentration
Meridian (AQSID 160010010)
Forecasted at 01/19/2022 Afternoon
Observation
Met Model Production
Bias Corrected Met Model Production
PM2.5(ug/m3)
35
30
25
20
15
10
5
01/12
01/13
01/14
01/15
01/16
01/17
01/18
01/19
01/20
Day


PM2.5 Daily AQI
Meridian (AQSID 160010010)
Forecasted at 01/19/2022 Afternoon
Observation
Met Model Production
Bias Corrected Met Model Production
PM2.5 AQI
0
5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
01/12
01/13
01/14
01/15
01/16
01/17
01/18
01/19
01/20
Day


PM2.5 Daily Forecast Bias
Meridian (AQSID 160010010)
PM2.5 Daily 24 Hour Average Concentration
Bias of Met Model Production
PM2.5(ug/m3)
15
10
5
0
−5
−10
01/12
01/13
01/14
01/15
01/16
01/17
01/18
PM2.5 Daily AQI
Bias of Met Model Production
PM2.5 AQI
25
20
15
10
5
0
−5
−10
−15
−20
01/12
01/13
01/14
01/15
01/16
01/17
01/18
Day


PM2.5 Hourly Concentration
Meridian (AQSID 160010010)
Forecasted at 01/19/2022 Afternoon
Observation
Met Model Production
PM2.5(ug/m3)
0
5
10
15
20
25
30
35
40
45
01/16_00
01/16_06
01/16_12
01/16_18
01/17_00
01/17_06
01/17_12
01/17_18
01/18_00
01/18_06
01/18_12
01/18_18
01/19_00
01/19_06
01/19_12
01/19_18
01/20_00
01/20_06
01/20_12
01/20_18
Hour (Local Standard Time)

Spreadsheet per Regional Office





Source : Internet

Neural Network

- Dense Neural Network
- 1D Convolutional Neural Network
- Recurrent Neural Network (LSTM)







Tree based Methods

- XGBoost
  - XGBoost stands for e**X**treme **G**radient **B**oosting
  - Tree built sequentially by minimizing the residue (error) of the previous tree
- Random Forest
- Boosted Random Forest







Precision
Recall
F1 Score

$$F = 2 \cdot \frac{\mathrm{precision} \cdot \mathrm{recall}}{\mathrm{precision} + \mathrm{recall}}$$

A measure that combines precision and recall is the harmonic mean of precision and recall, the traditional F-measure or balanced F-score

Source : Internet

Heidke Skill Score (HSS) and Hanssen-Kuiper Skill Score (KSS)

- HSS represents the accuracy of the model prediction compared with a reference forecast, which is from the random guess that is statistically independent of the observations.
- The range of the HSS is from -inf to 1. A negative value means a random guess is better, 0 means no skill, and 1 means a perfect score.
- KSS measures the ability to separate different categories. The range is from -1 to 1 where 0 means no skill, and 1 means a perfect score.

Violin Plot Explained

Present Distribution of Model Ensemble Member Forecasts







Figure 1. Common Components of Box Plot and Violin Plot. Total compensation for all academic ranks.



Source : Internet

NASA GEOS

GEOS Forecast and Reanalysis Products

GEOS Forward Processing (GEOS FP) NRT Analysis and Forecast

GEOS-Composition Forecast (GEOS-CF) NRT Forecast

Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) Reanalysis

ARSET - Introduction and Access to Global Air Quality Forecasting Data and Tools | NASA Applied Science

NASA GEOS-FP

GEOS FP

https://gmao.gsfc.nasa.gov/weather_prediction/

- GEOS FP analyses and forecasts support NASA field campaigns and provide a testbed for assimilation and forecast development
- Publicly available
- Includes weather, aerosols, and carbon monoxide (CO) on the same spatial scale
- State of the science forecast system – model physics or observing system updated every 6-12 months
  - Not suitable for trend analyses

AOD (550 nm)



https://svs.gsfc.nasa.gov/31100

- Meteorology used to drive chemistry models:
  - GEOSChem, Whole Atmosphere Community Climate Model (WACCM)
- When using FP meteorology fields to drive another model, must ensure your simulation does not span an update
  - GMAO NRT Product Page has updated details and dates

ARSET - Introduction and Access to Global Air Quality Forecasting Data and Tools | NASA Applied Science

NASA GEOS-CF

NASA Composition Forecasts (GEOS-CF)

GEOS Meteorology



GEOS-Chem Chemistry



GEOS-CF

- GEOS-Chem: Global chemistry transport model driven by GEOS meteorology
- 1-day simulation of the previous day using the analysis from FP-IT
  - Uses a **replay** technique to force the meteorology towards the FP-IT analysis
  - FP-IT is a 'frozen' version of FP used for satellite retrievals, similar to the version used to make MERRA-2.
- 5-day forecast
- Two aerosol schemes:
  - GOCART – Radiatively coupled to AGCM
  - GEOS-Chem – No feedbacks to model physics
- Full description in Keller et al., 2021

NASA GEOS MERRA-2

MERRA-2 Reanalysis

https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/

- The **M**odern-**E**ra **R**etrospective analysis for **R**esearch and **A**pplications version **2** (MERRA-2) provides data beginning in 1980 and runs a few weeks behind real-time.
- Long-term, model-based analyses of multiple datasets using a fixed assimilation system
- Includes meteorology, stratospheric ozone, and aerosols at the spatial resolution of a 0.5° × 0.66° (~50 km) grid.

Source: https://gmao.gsfc.nasa.gov/reanalysis/

ARSET - Introduction and Access to Global Air Quality Forecasting Data and Tools | NASA Applied Science

NASA GEOS Output

GEOS Output Quick Guide

	GEOS FP	GEOS-CF	MERRA-2
Type	Analysis + Forecast	Replay + Forecast	Reanalysis
Domain	Global	Global	Global
Spatial Resolution	Simulation: ~12 km Output: ~25 km (0.25°x0.312°)	~25 km (0.25°x0.312°)	~50km (0.5°x0.625°)
Temporal Resolution	2-D data: Hourly 3-D data: Every 3 h	15 min, Hourly	Hourly, Daily, Monthly
Vertical Levels	72 (near surface-0.1 hPa)	72 (near surface-0.1 hPa)	72 (near surface-0.1 hPa)
Output available	Analysis: 2014 – Present Forecast: ~20 days	Replay: 2018 – Present Forecast: 2019 – Present (aqc collection) ~14 days (all collections)	1980-Present
Initialization	Daily 10-day forecast at 00Z Daily 5-day forecast at 12Z	Daily 5-day forecast at 12Z	~1-2 months behind real time
Data Assimilation	Yes	No	Yes
File Specification Doc	https://gmao.gsfc.nasa.gov/pubs/docs/Lucchesi1203.pdf *	https://gmao.gsfc.nasa.gov/pubs/docs/Knowland1204.pdf *	https://gmao.gsfc.nasa.gov/pubs/docs/Bosilovich785.pdf *

* Find most current File Specification at https://gmao.gsfc.nasa.gov/pubs/office_notes.php

ARSET - Introduction and Access to Global Air Quality Forecasting Data and Tools | NASA Applied Science