

MOVES-AIRPACT4 Integration 2012 Update

June 7, 2012

Northwest Airquest MOVES Workgroup

AIRPACT4 Domain and Counties



 AIRPACT4 domain


States

-  California
-  Idaho
-  Montana
-  Nevada
-  Oregon
-  Utah
-  Washington
-  Wyoming
-  Canada

150

 Kilometers


**Expanded Modeling
Domain for AIRPACT4**



Why Switch Mobile Source Models?


- MOVES replaced MOBILE6.2 by EPA in March 2010
 - Required for regulatory purposes by March 2013
- MOVES model performance significantly improved compared to MOBILE6.2
 - WSU CO/NO_x study highlighted problem in PNW

2009 WSU Winter PM_{2.5} Boise Study




Source of base chart: Will Wallace, WSU, MOVES detail added by Idaho DEQ

Urban Arterials and Interstate dominate St. Lukes monitoring site



2009 WSU Winter PM_{2.5} Boise Study



Source of base chart: Will Wallace, WSU, MOVES detail added by Idaho DEQ

MOVES Workgroup Activities To-Date

- MOVES Workgroup formed June 2011
- Developed model integration plan
- Defined “County Groups” and “Reference Counties”
- Defined “Fuel Months” for summer and winter seasons
- Generated input files based on MOVES default data
- Test run SMOKE-MOVES using the “standard approach”
- Work in progress to develop Temperature profiles for each County Group
- Work in progress to setup SMOKE-MOVES in AIRPACT4 approach

Plan

Step 1: Establish the SMOKE-MOVES integration process in AIRPACT4 using mostly “national default” inputs.

Step 2: Refine Input files using local specific data for MOVES and SMOKE

Reference Counties

Purpose:

“...to reduce the computational burden running MOVES on every county in your modeling domain.”¹

¹User's Guide for the SMOKE-MOVES Integration Tool, July 14, 2010, USEPA OAQPS and UNC Institute for the Environment

What *IS* a County Group/ Reference County Anyway?

County Group


- A set of similar counties within a larger geographic area:
 - Fuel parameters
 - Fleet age distribution
 - Inspection and maintenance (I/M) programs

Reference County

- A county chosen to represent each County Group
- Chosen for their “averageness” with respect to the vehicle age distribution weighted by vehicle populations


AIRPACT4 County Fuel Groups

130 Kilometers




 AIRPACT4 domain boundary

Idaho




AIRPACT4 Oregon County Fuel Groups and Met Sites



130 Kilometers

AIRPACT4 Washington County Fuel Groups and Met Sites



Fuel Months

Concept:

“...is used to indicate when a particular set of fuel properties should be used in a MOVES simulation.”¹

and


“...similar to the reference county, the fuel month reduces the computational time of MOVES by using a single month to represent a set of months.”¹

¹User’s Guide for the SMOKE-MOVES Integration Tool, July 14, 2010, USEPA OAQPS and UNC Institute for the Environment

Fuel Month Designations

- Two Fuel Months chosen for MOVES-AIRPACT Integration
 - January
 - Represents Winter (November -- April)
 - July
 - Represents Summer (May -- October)

Basic SMOKE-MOVES Integration Tool Flowchart



SMOKE MOVES PROCESS CATEGORY


MOVES Lookup Table	Units	smokeProcID	Emissions Process
RatePerDistance (RPD)	Grams/mile	EXR CXR TIR BRK EPM EFL EFV	Running Exhaust Crankcase Running Exhaust Tire Wear Brake Wear On-road Evaporative Permeation On-road Evaporative Fuel Leaks On-road Evaporative Fuel Vapor Venting
RatePerVehicle (RPV)	Grams/vehicle/hour	EXS CXS EPM EFL CEI EXT	Start Exhaust Crankcase Start Exhaust Off-network Evaporative Permeation Off-network Evaporative Fuel Leaks Crankcase Extended Idle Exhaust Extended Idle Exhaust
RatePerProfile (RPP)	Grams/vehicle/hour	EFV	Off-network Evaporative Fuel Vapor Venting

Possible Approaches for AIRPACT-4 MOVES Integration


- Standard SMOKE Process Approach
 - Run MOVES on fly
 - No program changes for SMOKE process
 - Running time is not acceptable
 - Take ~26 days to generate lookup tables using 8 workers/CPU's for one day forecast
- Static Lookup Table Approach
 - Generate static lookup table in advance (one time deal, take long time to run)
 - Need to change SMOKE program and scripts
 - Shorten day to day process time significantly

Generate Lookup Table

Standard Way




AIRPACT4 Way




Historical Observed Meteorology

Historical Temperature Range will determine number of bins in Look-up tables.




Historical Observed Meteorology

Shape of Diurnal Temperature Profile affects the “Rate Per Profile” evaporative venting emissions – one set for each Reference County/Fuel Month Group




RatePerDistance (On-Network)

Standard Way




AIRPACT4 Way




RatePerVehicle and RatePerProfile (Off-Network)

Standard Way



AIRPACT4 Way




Test Run Result (Sunday, 4/1/2012)

RatePerDistance

Layer 1 BENZENE[1]

[1]=mgts_l_onroad.20120401.1.AIRPACT_04km.MOVES.ncf




April 2, 2012 00:00:00 UTC

Min (1, 1) = 0.000, Max (62, 199) = 0.002

Max. cells Layer: 1 BENZENE[1]

[1]=mgts_l_onroad.20120401.1.AIRPACT_04km.MOVES.ncf




■ (62, 199)

Test Run Result (Sunday, 4/1/2012)

RatePerVehicle

Layer 1 BENZENE[2]

[2]=mgts_l.offroad-rpv.20120401.1.AIRPACT_04km.MOVES.ncf




April 2, 2012 00:00:00 UTC

Min (1, 1) = 0.000, Max (62, 199) = 0.009

Max. cells Layer: 1 BENZENE[2]

[2]=mgts_l.offroad-rpv.20120401.1.AIRPACT_04km.MOVES.ncf




■ (278, 12)

Test Run Result (Sunday, 4/1/2012)

RatePerProfile

Layer 1 BENZENE[3]

[3]=mgts_offroad-rpp.20120401.1.AIRPACT_04km.MOVES.ncf




April 1, 2012 20:00:00 UTC

Min (1, 1) = 0.000E0, Max (62, 199) = 1.711E-4

Max. cells Layer: 1 BENZENE[3]

[3]=mgts_offroad-rpp.20120401.1.AIRPACT_04km.MOVES.ncf



(02, 199)

Test Run Running Time

Linux Machine, 2.3GHz CPU
Exclude lookup table generation

RatePerDistance		RatePerVehicle		RatePerProfile	
SMOKE Program	Running Time	SMOKE Program	Running Time	SMOKE Program	Running Time
SMKINVEN	< 1 minute	SMKINVEN	< 1 minute	SMKINVEN	< 1 minute
SPCMAT	< 1 minute	SPCMAT	< 1 minute	SPCMAT	< 1 minute
GRDMAT	~3 minutes	GRDMAT	~3 minutes	GRDMAT	~3 minutes
TEMPORAL	~2 minutes				
MOVESMRG	~150 minutes	MOVESMRG	~23 minutes	MOVESMRG	~3 minutes

Challenges in Current Approach

- MOVES doesn't contain default input data for Canada
- No SMOKE SAPRC99 speciation profile and cross reference files for SMOKE-MOVES integration
 - Only available for CB4 or CB5
- Run time issues
 - Generating one static lookup table, ~26 days run time
 - Daily run time ~ 2.7hrs (mostly MOVES Merge program ~2.5hrs)

Possible Speed-up Approaches to reduce daily run time

- Reduce look-up table size.
 - Change temp bin size from 5 deg to 10 or 20 deg
 - Reduce species to PM2.5 & ozone precursors
 - (remove GHG and toxics for now)
- Faster processor
- Wait for planned EPA speed-up for MOVES-Merge program in SMOKE

Up Next...

- Finalize setting up SMOKE-MOVES in AIRPACT4 approach
- Try to reduce daily run time
- Produce the specific local input for MOVES and SMOKE
- Generate new lookup tables and integrate them into daily process