

Working together for clean air



Puget Sound
Clean Air Agency

pscleanair.org



Enhancing our ability to track woodsmoke emission reductions

Levoglucosan Measurements

Acknowledgments

→ Neil Frank, USEPA, OAQPS

- Project Lead, Funding Lead

→ Prof. Chris Simpson, U of WA

→ DR. Gretchen Onstad, U of Wa

- Analysis, final report

→ PSCAA

- Filters and funding

→ Ecology

- Filters

Woodsmoke is not just a Region 10 problem

- Woodsmoke drives high PM 2.5 levels in winter and is the chief contributor to non-attainment**
- Measuring OC via speciation network is expensive and to some extent confounded by other emission sources**
- Recent advances in analysis work suggest that Levoglucosan may be a viable marker for wood smoke**
- Existing filter archives can be used to establish baselines**


Levoglucosan (LG) is an indicator of Residential Wood Combustion PM2.5

→ Puget Sound Study (2004-2007 filters)

- 300 filters at 2 urban and 3 suburban/rural sites
- Ethyl acetate extraction procedure coupled with derivatization and GC/MS analysis using selected ion monitoring (SIM)

→ LG can also be measured with water extracts using HPLC-PAD

- This approach permits other useful analyses



See Anast and Simpson (2008)

LG% of PM_{2.5} is highest in Darrington

LG varies with local RWC contribution to PM_{2.5}




Figure 7: Associations between levoglucosan and PM mass at five monitoring sites in the Puget Sound Airshed.

Table 4: Empirically derived scaling factors indicating the relationship between levoglucosan and PM_{2.5} mass for woodsmoke dominated periods and locations in the Puget Sound Airshed.

	Average % levoglucosan per µg PM _{2.5}	95% confidence interval
Darrington	13.9	12.7-15.2
South Tacoma	9.6	8.6-10.5
Marysville	10.0	9.1-10.7


Measurement of LG in archived FRM filters

- LG is reliably measured in archived FRM filters
- Good agreement is observed between organic solvent extraction/GCMS and aqueous extraction/HPLC-PAD methods.

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

LG and source apportionment

- LG is strongly correlated with the putative “woodsmoke source” factor in PMF-derived source apportionment models, and is not correlated with other factors in the model.
- This provides confidence that the woodsmoke source factor has been correctly identified in the model





LG and source apportionment

- When LG is explicitly included as a variable in the PMF model, a more accurate apportionment is achieved.
- Based on this model, LG comprises ~ 9% of woodsmoke-derived PM mass. This is consistent with values previously reported in a number of source testing studies.


QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

Spatial distribution of woodsmoke concentrations

- Measurement of LG indicates that temporal and spatial variation in woodsmoke-PM is greater than variation in total $PM_{2.5}$ concentrations.
- This information assists local agencies to identify areas where woodsmoke drives air quality concerns, and thence to implement appropriate, targeted interventions.



LG to woodsmoke PM scaling factors



Scaling factors obtained from ambient samples

- A wide range of scaling factors have been reported in the literature from source-testing studies, scaling factors calculated from ambient measurements show less variability.
- This indicates that it is feasible to use measurements of LG to reliably estimate the woodsmoke contribution to ambient PM_{2.5}.
- Data from the Darrington site (above) suggest that significant regional differences exist in the LG to woodsmoke PM scaling factors..

Conclusions- U of WA

- **LG can be reliably measured on archived FRM (Teflon) filters**
Two analytical methods have been validated for this purpose, and have been shown to give excellent agreement:
 - **Organic extraction followed by GC/MS analysis**
 - **Aqueous extraction followed by HPLC (ion exchange) with PAD detection**
- **Measurement of LG allows agencies to identify seasons and locations where woodsmoke drives PM concentrations (and potential NAAQS violations). This in-turn permits implementation of appropriately targeted interventions to reduce PM concentrations.**
- **Within a geographic area the scaling factors between LG and ambient woodsmoke-derived PM_{2.5} mass is relatively precise. Therefore, woodsmoke-derived PM_{2.5} mass can be estimated with acceptable precision from a limited number of measurements of LG.**

LG and other analyses using archived Teflon filters to track emission reductions

→ Percent LG will help determine relative contribution of RWC

- As RWC decreases, LG will also decrease

- ▲ If % LG decreases, then RWC is responsible for reduced PM_{2.5}

- ▲ If %LG stays the same, then other sources may also be decreasing

- ▲ If %LG increases, than other sources are responsible

Recommendation

- **NWAIRQUEST should further evaluate**
- **If warranted seek new funding for adding routine analysis of Levoglucosan into Region 10 monitoring networks.**
- **Include in SIP/attainment plans to track emission reduction and effectiveness of wood stove change out programs**

LG and other analyses using archived Teflon filters to track emission reductions

→ For further information, see

- Zhang and Hecobian -- http://www.atmos-chem-phys-discuss.net/papers_in_open_discussion.html
- Frank [Levoglucosan and Water Soluble Organic Carbon from PM2.5 Teflon Filters \(PDF\)](#), Natl Air Monitoring Conference (2009)
<http://www.epa.gov/ttnamti1/files/ambient/2009conference/frank.pdf>