

Idaho Department of Environmental Quality

Statistical analysis of AIRPACT ozone forecasts
in the Treasure Valley of southwest Idaho

Kimi Smith, Technical Services Division



Motivation

- In 2019, AIRPACT added no skill when forecasting daily AQI compared to the IDEQ regional forecasters
- Ozone is a pollutant of concern in SW Idaho



Table A3. 2017–2019 ozone preliminary design values.

| Site | County/ AQS ID | 4th-Highest Daily Maximum 8-Hour Average (ppm) | | | 3-Year Design Value (ppm) |
|-------------------------|-------------------|---|-------------|-------------|------------------------------|
| | | 2017 | 2018 | 2019 | |
| Boise—White Pine | Ada 160010017 | 0.076/0.068 | 0.068/0.064 | 0.052/0.052 | 0.065/0.061 |
| Meridian— St. Luke's | Ada 160010010 | 0.071/0.069 | 0.067/0.066 | 0.057/0.057 | 0.065/0.064 |

Notes: A monitor violates the 8-hour ozone National Ambient Air Quality Standard if the 3-year average of the annual 4th-highest daily maximum average exceeds 0.070 parts per million (ppm). Concentration data are denoted with/without exceptional event data included. AQS = air quality system; ID = identification

Data

- From the AIRPACT website: downloaded modeled hourly ozone concentrations and AIRNow observations at the St. Luke's site in Meridian for 2017, 2018, 2019, 2020 (Aug.)



Methods

- Completed a statistical analysis including:
 - Categorical and error statistics: false alarm rate (FAR), probability of detection (POD), critical success index (CSI), root mean square error (RMSE), scatter index (SI), mean absolute error (MAE)
 - Forecast verification: two-category forecast table

Categorical and Error Statistics

- Selected AQI > 50 as the threshold because IDEQ forecasters are most accurate at forecasting AQI below that threshold
- Accuracy drops off considerably at AQI values above 50 – most critical for public health
- Is AIRPACT helpful for forecasting values above AQI 50?
- AQI 50 = 54 ppb of ozone

| Year-Long (AQI > 50) | | | |
|----------------------|-------|-------|-------|
| Statistic | 2017 | 2018 | 2019 |
| FAR | 3.39 | 1.34 | 13.95 |
| POD | 45.71 | 21.67 | 71.43 |
| CSI | 40.00 | 20.31 | 16.39 |

| Ozone Season (April - September) | | | |
|----------------------------------|-------|-------|------|
| Statistic | 2017 | 2018 | 2019 |
| RMSE | 12.89 | 8.03 | 8.42 |
| Scatter Index | 0.25 | 0.16 | 0.19 |
| MAE | 7.84 | 6.32 | 6.48 |
| FCST-OBS avg | -1.60 | -4.08 | 4.54 |

Forecast Verification


| TWO-CATEGORY FORECAST TABLE | | Observed | | |
|--------------------------------|-------|-------------|-------------------------|--------------|
| | | Yes | No | Total |
| Forecast | Yes | Hit | False Alarm | Forecast Yes |
| | No | Miss | Correct Negative | Forecast No |
| | Total | Obs. Yes | Obs. No | Total |

| 2018 (AQI > 50) | | Observed | | |
|------------------------|-------|-----------|------------|-------|
| | | Yes | No | Total |
| Forecast | Yes | 3 | 4 | 7 |
| | No | 47 | 294 | 341 |
| | Total | 50 | 298 | 348 |

| 2019 (AQI > 50) | | Observed | | |
|------------------------|-------|-----------|------------|-------|
| | | Yes | No | Total |
| Forecast | Yes | 10 | 47 | 57 |
| | No | 4 | 290 | 294 |
| | Total | 14 | 337 | 351 |


| 2017 (AQI > 50) | | Observed | | |
|------------------------|-------|-----------|------------|-------|
| | | Yes | No | Total |
| Forecast | Yes | 32 | 10 | 42 |
| | No | 38 | 285 | 323 |
| | Total | 70 | 295 | 365 |

False Alarm Days by Month




April-May Comparison

Difference between AIRPACT forecast ozone and observed ozone (ppb)



April-May Comparison

Average of the difference between the AIRPACT forecast ozone and observed ozone over April-May (ppb)



2020 Forecast Verification (So Far)

- Data from January through August

| 2020 (AQI > 50) | | Observed | | |
|-----------------|-------|-----------|------------|-------|
| | | Yes | No | Total |
| Forecast | Yes | 15 | 35 | 50 |
| | No | 10 | 184 | 194 |
| | Total | 25 | 219 | 244 |

| 2019 (AQI > 50) | | Observed | | |
|-----------------|-------|----------|------------|-------|
| | | Yes | No | Total |
| Forecast | Yes | 9 | 45 | 54 |
| | No | 4 | 175 | 179 |
| | Total | 13 | 220 | 233 |

Summary

- AIRPACT over-predicted ozone in 2019 and 2020 compared to 2017 and 2018
 - Most frequent over-prediction occurred in the spring months (March, April, May)
- AIRPACT doesn't appear to improve daily AQI forecasts in SW ID when the AQI is above 50

Potential Causes?

- Meteorology-related?
 - Slightly cooler and much wetter in spring 2019 compared to spring 2017/2018; spring 2020 similar to 2017/2018 – we plan to look into the meteorology further
- Biomass burning-related?
- Possible change(s) in AIRPACT?

Future Work

- Compare AIRPACT performance with observations at other sites in/around Idaho
- Evaluate AIRPACT forecasts for other pollutants (PM)

Thank you!

- We welcome any thoughts and/or feedback!

Contact

Kimi Smith
kimi.smith@deq.idaho.gov



Statistic Calculations


- **False alarm rate (FAR)**
 - Percent of time the event was forecast but did not occur
 - $\text{False Alarms} / (\text{False Alarms} + \text{Hits}) * 100$
- **Probability of detection (POD) / hit rate**
 - Ability of model to predict events
 - $\text{Hits} / (\text{Hits} + \text{Misses}) * 100$
- **Critical success index (CSI) / threat score**
 - How well events were predicted
 - $\text{Hits} / (\text{Hits} + \text{False Alarms} + \text{Misses}) * 100$

Statistic Calculations


- **Root mean square error (RMSE)**
 - Average magnitude of forecast errors; puts more weight on larger errors
 - $\text{SqRt}(\text{SumSq}(\text{Forecast} - \text{Observed})/\text{Total})$
- **Scatter index (SI)**
 - Expected error within one standard deviation; if above 1 (or 100%), error is more than one deviation
 - $\text{RMSE} / \text{Average}(\text{Total})$
- **Mean absolute error (MAE)**
 - Average magnitude of forecast errors; calculated on the absolute value of the differences
 - $\text{Average}(\text{AbsoluteValue}(\text{Forecast} - \text{Observed}))$

April-May Comparison

Average of the absolute difference between the AIRPACT forecast ozone and observed ozone (MAE) over April-May (ppb)



2017-2020 FCST-OBS & 7-day MA



2017-2020 14-day & 30-day MA

